

Supplementary References (Section: Bioinformatics and statistical analysis of HT-NGS data)

- S1. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet Journal* 2011; 17(1), 10-12.
- S2. Kozomara A, Griffiths-Jones S. miRBase: annotating high confidence microRNAs using deep sequencing data. *Nucleic Acids Res* 2014; 42: D68-D73.
- S3. Ashurst JL, Chen CK, Gilbert JGR, Jekosch K, Keenan S, Meidl P, *et al.* The Vertebrate Genome Annotation (Vega) database. *Nucleic Acids Res* 2005; 33: D459-D465.
- S4. Aken BL, Ayling S, Barrell D, Clarke L, Curwen V, Fairley S, *et al.* The Ensembl gene annotation system. *Database* 2016, baw093. doi: 10.1093/database/baw093.
- S5. O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, *et al.* Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res* 2016; 44(D1): D733-45.
- S6. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol* 2009; 10: R25.
- S7. Friedländer MR, Chen W, Adamidi C, Maaskola J, Einspanier R, Knespel S, *et al.* Discovering microRNAs from deep sequencing data using miRDeep, *Nature Biotechnol* 2008; 26: 407-15.
- S8. Langmead B, Salzberg S. Fast gapped-read alignment with Bowtie 2. *Nature Methods* 2012; 9: 357-59.
- S9. Bo Li and Colin N Dewey. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* 2011; 12: 323 DOI:10.1186/1471-2105-12-323.
- S10. Robinson MD, Oshlack A. A scaling normalization method for differential expression analysis of RNA-seq data. *Genome Biol* 2010; 11: R25.
- S11. Zhou X, Lindsay H, Robinson MD. Robustly detecting differential expression in RNA sequencing data using observation weights. *Nucleic Acids Res* 2014; 42(11): e91.
- S12. Love MI, Huber W and Anders S. Moderated estimation of fold change and dispersion for RNA-Seq data with DESeq2. *Genome Biol* 2014; 15(12): 550.
- S13. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, *et al.* Limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Research* 2014; 43(7): e47.
- S14. Law C, Chen Y, Shi W, Smith GK. voom: precision weights unlock linear model analysis tools for RNA-seq read counts. *Genome Biol* 2014; 15: R29.

- S15.** Zhang Y, Verbeek FJ. Comparison and integration of target prediction algorithms for microRNA studies. *J Integr Bioinform* 2010; 7(3). doi: 10.2390/biecoll-jib-2010-127.
- S16.** Grimson A, Farh KK, Johnston WK, Garrett-Engele P, Lim LP, Bartel DP. MicroRNA targeting specificity in mammals: determinants beyond seed pairing. *Mol Cell* 2007; 27: 91–105.
- S17.** Baek D, Villen J, Shin C, Camargo FD, Gygi SP, Bartel DP. The impact of microRNAs on protein output. *Nature* 2008; 455: 64-71.